

# Anti- human E-Cadherin Mouse Monoclonal Primary Antibody

Clone: OTI1F3

**IVD**

**REF** CE00075

## CATALOG NUMBER

C0075MA01-MA 0.1 mL  
C0075MA05-MA 0.5 mL  
C0075MA10-MA 1.0 mL

## ENGLISH

### Intended use

Anti- human E-Cadherin (Clone: OTI1F3) Mouse Monoclonal Primary Antibody is intended for detection of E-Cadherin protein expression in frozen or formalin fixed human tissues and cells. The clinical interpretation of any positive staining or its absence should be complemented by morphological and histological studies with proper controls. Evaluations should be made within the context of the patient's clinical history and other diagnostic tests by a qualified pathologist. The antibody is intended for *in vitro* diagnostic (IVD) use.

### Background

This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a calcium dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function is thought to contribute to progression in cancer by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. Identified transcript variants arise from mutation at consensus splice sites. [provided by RefSeq, Jul 2008].

Alternative names: Arc-1; CD324; CDHE; ECAD; LCAM; UVO

### Reagent provided

Anti-human E-Cadherin Mouse Monoclonal Primary Antibody (Clone: OTI1F3) is provided in liquid form in 20mM Sodium phosphate, 150mM Sodium chloride, 0.2% BSA, 0.09% Sodium azide, pH 7.4. The isotype of the antibody is IgG1,k. The protein concentration is approximately 1.0 +/- 0.05 mg/mL.

For immunohistochemistry, the primary antibody may be used at a working dilution of 1:100 – 1:200 for formalin-fixed, paraffin-embedded human tissue. It can be dependent upon the detection system used. These are guidelines only, and optimal dilutions should be determined by the individual laboratory.

### Immunogen

Full length human recombinant protein of human CDH1 (NP\_004351) produced in HEK293T cell.

## Specificity

The specificity of the anti- human E-Cadherin Mouse Monoclonal Primary Antibody was established on human kidney. The anti-human E-Cadherin presented no staining on human brain and positive staining on human kidney using immunohistochemical (IHC) test methods.

## Materials Required but Not Supplied

Antibody diluent, HIER solution, Antibody detection kits, Chromogen, Staining reagents, negative and positive tissue control slides are not included.

## Precautions

1. For use by trained professionals only.
2. This product contains sodium azide ( $\text{NaN}_3$ ), a chemical highly toxic in pure form. At product concentrations, though not classified as hazardous,  $\text{NaN}_3$  may react with lead and copper plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent metal azide build-up in plumbing.
3. Wear appropriate Personal Protective Equipment to avoid contact with eyes and skin.
4. Unused reagents should be disposed of according to local, State, and Federal regulations.

## Storage

Store at 2-8°C. Do not use the product past the expiration date indicated on the label. If reagents are stored under any other conditions, the end user must verify the acceptability of those conditions. There are no obvious signs to indicate instability of this product therefore, positive and negative controls should be run simultaneously with patient specimens.

## Specimen Preparation

### Paraffin Sections

Anti- human E-Cadherin Mouse Monoclonal Primary Antibody can be used on formalin-fixed, paraffin-embedded tissue sections at a working dilution of 1:100 to 1:200. Anti- human E-Cadherin Mouse Monoclonal Primary Antibody (Clone: OT11F3) working dilution requires heat induced epitope retrieval (HIER) for 3 minutes using pressure chamber at 110C for staining. We recommend using HIER Citrate solution pH 6.0, which showed optimal staining of anti-E-Cadherin antibody at a dilution of 1:100 on human kidney. The dilutions are estimates; the actual staining results may vary due to reagents and detection protocols used. Validation of antibody performance and final protocol are the responsibility of the end user.

## Staining procedure

### Manual Staining Procedure

1. Deparaffinize slides.
2. Submerge slides in peroxidase quenching solution for ~10 minutes, then rinse 2x with dH<sub>2</sub>O.
3. Heat Induced Epitope Retrieval is required for this antibody; Citrate solution, pH 6.0 at 110C for 3minutes.
4. Allow slides to cool down from step 3, rinse with distilled water, wash with PBS-T 3 times, 2 minutes each.
5. Apply serum blocking solution.[Optional]
6. Apply primary antibody and incubate for 30-60 minutes at room temperature. After incubation wash with PBS-T 3 times, 2 minutes each.
7. Apply secondary antibody and incubate according to the data sheet of the detection system. Wash with PBS-T 3 times, 2 minutes each.
8. Apply enzyme conjugate and incubate according to data sheet of detection system. Wash with PBS-T 3 times, 2 minutes each.
9. Apply chromogen and incubate 5-10 minutes and rinse with distilled water.

## Staining interpretation

The cellular staining pattern for Anti- human E-Cadherin Mouse Monoclonal Primary Antibody is membranous and cytoplasmic.

## Performance Characteristics

### Predicted Staining in Normal Tissue/Cells

Human brain was shown to be negative for this antibody.

### Predictive Staining in Tumor

Anti- human E-Cadherin Mouse Monoclonal (Clone: OT11F3) produced membranous and cytoplasmic staining when screened on human kidney.

## Contact Information



SDIX LLC  
111 Pencader Drive  
Newark, Delaware 19702  
USA  
+1 302 456 6789  
+1 800 544 8881(USA)  
[www.SDIX.com](http://www.SDIX.com)

## Product Complaint and/or Technical Support

[techsupport@origene.com](mailto:techsupport@origene.com)  
+1 301 340 3188 (prompt 2)

## Authorized Representative

Colin LeGood  
Barnes Wallis House, 25 Barnes Wallis Road  
Segensworth East, Hampshire PO15 5TT UK  
Tel +44 (0) 1489 898640